

### **Publications using RMD mutant lines:**

1. Li F, Zhang M, Guo K, Hu Z, Zhang R, Feng Y, Yi X, Zou W, Wang L, Wu C, Tian J, Lu T, Xie G, Peng L. High-level hemicellulosic arabinose predominately affects lignocellulose crystallinity for genetically enhancing both plant lodging resistance and biomass enzymatic digestibility in rice mutants. *Plant Biotechnol J*. 2015 May;13(4):514-525.
2. Chang Y, Gong L, Yuan W, Li X, Chen G, Li X, Zhang Q, Wu C. Replication Protein A (RPA1a) is required for meiotic and somatic DNA repair but is dispensable for DNA replication and homologous recombination in rice. *Plant Physiol*, 2009, 151: 2162-2173.
3. Chu H, Liang W, Li J, Hong F, Wu Y, Wang L, Wang J, Wu P, Liu C, Zhang Q, Xu J, Zhang D. A CLE-WOX signaling module regulates root meristem maintenance and vascular tissue development in rice. *J Exp Bot*, 2013, 64: 5359-5369.
4. Dai X, You C, Chen G, Li X, Zhang Q, Wu C. OsBC1L4 encodes a COBRA-like protein that affects cellulose synthesis in rice. *Plant Mol Biol*, 2011, 75: 333-345.
5. Dai X, You C, Wang L, Chen G, Zhang Q, Wu C. Molecular characterization, expression pattern, and function analysis of the OsBC1L family in rice. *Plant Mol Biol*, 2009, 71: 469-481.
6. Deng H, Liu H, Li X, Xiao J, Wang S. A CCCH-Type zinc finger nucleic acid-binding protein quantitatively confers resistance against rice bacterial blight disease. *Plant Physiol*, 2012, 158: 876-889.
7. Ding X, Cao Y, Huang L, Zhao J, Xu C, Li X, Wang S. Activation of the indole-3-acetic acid-amido synthetase GH3-8 suppresses expansion expression and promotes salicylate- and jasmonate-independent basal immunity in rice. *Plant Cell*, 2008, 20: 228-240.
8. Du H, Liu L, You L, Yang M, He Y, Li X, Xiong L. Characterization of an inositol 1,3,4-trisphosphate 5/6-kinase gene that is essential for drought and salt stress responses in rice. *Plant Mol Biol*, 2011, 77: 547-563.
9. Du H, Wang N, Cui F, Li X, Xiao J, Xiong L. Characterization of the carotene hydroxylase gene DSM2 conferring drought and oxidative stress resistance by increasing xanthophylls and abscisic acid synthesis in rice. *Plant Physiol*, 2010, 154:1304-1318.
10. Gao X, Chen Z, Zhang J, Li X, Chen G, Li X, Wu C. OsLIS-L1 encoding a lissencephaly type-1-like protein with WD40 repeats is required for plant height and male gametophyte formation in rice. *Planta*, 2012, 235: 713-727.
11. Hu Y, Liu D, Zhong X, Zhang C, Zhang Q, Zhou D. CHD3 protein recognizes and regulates

methylated histone H3 lysines 4 and 27 over a subset of targets in the rice genome. *Proc Natl Acad Sci USA*, 2012, 109: 5773-5778.

12. Jiang Y, Bao L, Jeong SY, Kim SK, Xu C, Li X, Zhang Q. XIAO is involved in the control of organ size by contributing to the regulation of signaling and homeostasis of brassinosteroids and cell cycling in rice. *Plant J*, 2012, 70: 398-408.
13. Liang D, Wu C, Li C, Xu C, Zhang J, Kilian A, Li X, Zhang Q, Xiong L. Establishment of a patterned GAL4-VP16 Transactivation system for discovering gene function in rice. *Plant J*, 2006, 46: 1059-1072
14. Li A, Zhang Y, Wu X, Tang W, Wu R, Dai Z, Liu G, Zhang H, Wu C, Chen G, Pan X. DH1, a LOB domain-like protein required for glume formation in rice. *Plant Mol Biol*, 2008, 66: 491-502.
15. Li J, Chu H, Zhang Y, Mou T, Wu C, Zhang Q, Xu J. The Rice HGW gene encodes a ubiquitin-associated (UBA) domain protein that regulates heading date and grain weight. *PLoS One*, 2012, 7: e34231.
16. Li X, Chang Y, Xin X, Zhu C, Li X, Higgins JD, Wu C. Replication protein A2c coupled with replication protein A1c regulates crossover formation during meiosis in rice. *Plant Cell*, 2013, 25:3885-3899.
17. Li X, Gao X, Wei Y, Deng L, Ouyang Y, Chen G, Li X, Zhang Q, Wu C. Rice APOPTOSIS INHIBITOR5 coupled with two DEAD-box adenosine 5'-Triphosphate-dependent RNA helicases regulates tapetum degeneration. *Plant Cell*, 2011, 23: 1416-1434.
18. Li X, Yang Y, Yao J, Chen G, Li X, Zhang Q, Wu C. FLEXIBLE CULM 1 encoding a cinnamyl-alcohol dehydrogenase controls culm mechanical strength in rice. *Plant Mol Biol* , 2009, 69: 685-697.
19. Long T, Guo D, He D, Shen W, Li X. The tRNA 3'-end processing enzyme tRNase Z2 contributes to chloroplast biogenesis in rice. *JIPB*, 2013, 55: 1104-1118.
20. Ning J, Li X, Hicks L, Xiong L. A Raf-Like MAPKKK gene DSM1 mediates drought resistance through reactive oxygen species scavenging in rice. *Plant Physiol*, 2010, 152: 876-890.
21. Ning J, Zhang B, Wang N, Zhou Y, Xiong L. Increased Leaf Angle1, a Raf-Like MAPKKK that interacts with a nuclear protein family, regulates mechanical tissue formation in the lamina joint of rice. *Plant Cell*, 2012, 23: 4334-4347.
22. Qi Z, Xiong L. Characterization of a purine permease family gene. *JIPB*, 2013, 55:

1119-1135.

23. Sang X, Li Y, Luo Z, Ren D, Fang L, Wang N, Zhao F, Ling Y, Yang Z, Liu Y, He G. (). CHIMERIC FLORAL ORGANS1, encoding a monocot-specific MADS box protein, regulates floral organ identity in rice. *Plant Physiol*, 2012, 160: 788-807.
24. Sasaki A, Yamaji N, Yokosho K, Ma JF. Nramp5 is a major transporter responsible for manganese and cadmium uptake in rice. *Plant Cell*, 2012, 24: 2155-2167.
25. Shen X, Liu H, Yuan B, Li X, Xu C, Wang S. OsEDR1 negatively regulates rice bacterial resistance via activation of ethylene biosynthesis. *Plant, Cell & Environ*, 2011, 34: 179-191.
26. Sun Q, Zhou D. Rice jmjC domain-containing gene JMJ706 encodes H3K9 demethylase required for floral organ development. *Proc Natl Acad Sci USA*, 2008, 105: 13679-13684.
27. Wu C, You C, Li C, Long T, Chen G, Byrne ME, Zhang Q. RID1, encoding a Cys2/His2-type zinc finger transcription factor, acts as a master switch from vegetative to floral development in rice. *Proc Natl Acad Sci USA*, 2008, 105: 12915-12920.
28. Wuriyanghan H, Zhang B, Cao W, Ma B, Lei G, Liu Y, Wei W, Wu H, Chen L, Chen H, Cao Y, He S, Zhang W, Wang X, Chen S, Zhang J. The ethylene receptor ETR2 delays floral transition and affects starch accumulation in rice. *Plant Cell*, 2009, 21: 1473-1494.
29. Xiang Y, Tang N, Du H, Ye H, Xiong L. Characterization of OsbZIP23 as a key player of the basic leucine zipper transcription factor family for conferring abscisic acid sensitivity and salinity and drought tolerance in rice. *Plant Physiol*, 2008, 148: 1938-1952.
30. Xu M, Galhano R, Wiemann P, Bueno E, Tiernan M, Wu W, Chung I, Gershenzon J, Tudzynski B, Sesma A, Peters R.J. Genetic evidence for natural product-mediated plant-plant allelopathy in rice (*Oryza sativa*). *New Phytol*, 2012, 193: 570-575.
31. Xu Y, McCouch S.R, Zhang Q. How can we use genomics to improve cereals with rice as a reference genome? *Plant Mol Biol*, 2005, 59: 7-26.
32. Xue-biao, P.C.L.A. Morphogenesis, Anatomical Observation and Primary Genetic Analysis of a Multi-glume Floral Organ Mutant in Rice. *Rice Science*, 2006, 13: 227-233.
33. Yang Y, Peng Q, Chen G, Li X., Wu C. OsELF3 is involved in circadian clock regulation for promoting flowering under long-day conditions in rice. *Mol Plant*, 2013, 6: 202-215.
34. You C, Dai X, Li X, Wang L, Chen G, Xiao J, Wu C. Molecular characterization, expression pattern, and functional analysis of the OsIRL gene family encoding intracellular Ras-group-related LRR proteins in rice. *Plant Mol Biol*, 2010, 74: 617-629.

35. Yuan B, Shen X, Li X, Xu C, Wang S. Mitogen-activated protein kinase OsMPK6 negatively regulates rice disease resistance to bacterial pathogens. *Planta*, 2007, 226: 953-960.
36. Yuan J, Chen D, Ren Y, Zhang X, Zhao J. Characteristic and expression analysis of a metallothionein gene, OsMT2b, down-regulated by cytokinin suggests functions in root development and seed embryo germination of rice. *Plant Physiol*, 2008, 146: 1637-1650.
37. Yuan W, Li X, Chang Y, Wen R, Chen G, Zhang Q, Wu C. Mutation of the rice gene. *The Plant J*, 2009, 59: 303-315.
38. Zhang C, Xu Y, Guo S, Zhu J, Huan Q, Liu H, Wang L, Luo G, Wang X, Chong K. Dynamics of brassinosteroid response modulated by negative regulator LIC in rice. *PLoS Genetics*, 2012, 8: e1002686.
39. Zhang C, Yuan W, Zhang Q. RPL1, a gene involved in epigenetic processes regulates phenotypic plasticity in rice. *Mol Plant*, 2012, 5: 482-493.
40. Zheng L, Yamaji N, Yokosho K, Ma JF. YSL16 is a phloem-localized transporter of the copper-nicotianamine complex that is responsible for copper distribution in rice. *Plant Cell*, 2012, 24: 3767-3782.
41. Zhu X, Xiong L. Putative megaenzyme DWA1 plays essential roles in drought resistance by regulating stress-induced wax deposition in rice. *Proc Natl Acad Sci USA*, 2013, 110: 17790-17795.
42. Zhang J, Guo D, Chang Y, You C, Li X, Dai X, Weng Q, Zhang J, Chen G, Li X, Liu H, Han B, Zhang Q, Wu C. Non-random distribution of T-DNA insertions at various levels of the genome hierarchy as revealed by analyzing 13,804 T-DNA flanking sequences from an enhancer-trap mutant library. *Plant J*, 2007, 49: 947-959
43. 何宗顺, 李雪梅, 吴昌银. 水稻穗大小决定基因 PS1 的遗传分析及克隆. *分子植物育种*, 2012, 10: 380-387.
44. 李爱宏, 武茹, 张亚芳, 汤雯, 吴昌银, 潘学彪. 一个水稻多重颖壳突变体的形态学观察及初步遗传分析. *中国水稻科学*, 2006, 20: 348-354.
45. 李雪梅, 何宗顺, 余四斌, 陈国兴, 吴昌银. 水稻穗形态建成基因 PMM1 的遗传分析与初步定位. *华中农业大学学报*, 2012, 31: 10-15.
46. 刘辉, 赵竹青, 陈涛. 水稻 T-DNA 插入突变体库筛选及其突变特征研究. *江苏农业科学*, 2007, 1: 1-3.
47. 刘辉, 赵竹青. 水稻 T-DNA 插入氮营养缺陷型突变体的初步筛选. *华中农业大学学报*, 2008, 27: 1-6.

48. 马玉银, 张亚芳, 潘存红, 李爱宏, 吴旭江, 左示敏, 陈宗祥, 吴昌银, 潘学彪. 水稻小粒矮突变性状与 T-DNA 插入标签的共分离检测. 中国水稻科学, 2008, 22: 571-577.
49. 裴荣, 陆展华, 姚家玲. 批量筛选水稻 T-DNA 插入突变体库获得生殖发育相关突变体. 华中农业大学学报, 2012, 31: 133-138.